arXiv:1402.2057v2 [cond-mat.str-el] 27 May 2014

Robustness of complex many-body networks: Novel perspective in 2D metal-insulator transition

Chung-Pin Chou
Beijing Computational Science Research Center, Beijing 100084, China

We present a novel theoretical framework established by complex network analysis for understanding the phase transition beyond the Landau symmetry breaking paradigm. In this paper we take a two-dimensional metal-insulator transition driven by electron correlations for example. Passing through the transition point, we find a hidden symmetry broken in the network space, which is invisible in real space. This symmetry is nothing but a kind of robustness of the network to random failures. We then show that a network quantity, small-worldness, is capable of identifying the phase transition with/without any symmetry breaking in the real space and behaving as a new order parameter in the network space. We demonstrate that whether or not the symmetry is broken in real space a variety of phase transitions in condensed matters can be characterized by the hidden symmetry breaking in the weighted network, that is to say, a decline in network robustness.

PACS numbers: 64.60.aq, 64.60.A-, 71.10.Fd, 71.30.+h

I. INTRODUCTION

Phases of matter are conventionally distinguished by using Landau's approach, which characterizes phases in terms of underlying symmetries that are spontaneously broken. The information we need to understand phase transitions is encoded in appropriate correlation functions, e.g., the correlation length would diverge close to a quantum critical point. Particularly, the low-lying excitations and the long-distance behavior of the correlations near the critical phase are believed to be well described by a quantum field theory. A major problem is, however, that in some cases it is unclear how to extract important information from the correlation functions if these phases do not break any symmetry, such as a nonmagnetic Mott transition [1].

Owing to the breakthrough in the control of ultracold atoms on optical lattices, Mott's original ideas about metal-insulator transitions have been realized in recent years [2–4]. His brilliant idea is well known as the Mott transition that a metal-insulator transition is driven by electron correlations. Although Mott insulators are usually related to antiferromagnetism, the Mott transition can also take place in the absence of long-range magnetic order. In the past years, much progress has been made from both theoretical and experimental sides in understanding Mott transitions [5]. A difficulty is the absence of the order parameter to characterize the critical behavior of the observables for the Mott transition not accompanied by the onset of antiferromagnetic order.

There are two earlier concepts as to what should be the order parameter to describe the physics around the Mott transition point from the metallic side. One is based on the disappearance of the Fermi liquid quasiparticles introduced by Brinkman and Rice [6]. However, the Brinkman-Rice transition actually exists only in infinite dimensions. Notice that the Mott transition still can be characterized by the charge stiffness, which is the response to a twist, while no symmetry is broken here [7]. The other concept is that the metal-insulator transition can be viewed as a condensation of doubly occupied sites (doublons), namely, the reduction of the number of carriers in the insulating side [8]. For simplicity, we will use the latter to study the Mott transition, and also assume the trial wave function with only nearest-neighbor attractive correlations between a doublon and an empty site (holon) in variational calculations.

In this paper, we propose the weighted network constructed by the correlation between a doublon and a holon in the two-dimensional (2D) Hubbard model on a square lattice. Complex network theory has become one of the most powerful frameworks for understanding the network structures of many real systems [9–13]. According to graph theory, the elements of the system often are called nodes and the relationships between them, which a weight is associated with, are called links. The weight of the network link contains useful information about topologies of the weighted network [14]. Decades ago, this unnoticed idea constructing a weighted network from condensed matters had been proposed in quantum Hall systems [15]. However, we here define the link weight carrying important information as doublon-holon binding correlations in the Hubbard model.

In Fig.1, one can see the network topologies corresponding to the metallic and insulating regimes (see the numerical details in Sec.II). The weighted network of the insulating state exhibits that there are only few links with the strongest weight called "highways" in the network, which looks very different from the metallic state. This observation reminds us of a well-known fact in real-world networks that heterogeneous



FIG. 1: Network representations of the 2D Hubbard model at interaction strength (a) $U = 0$ and (b) $U = 12$. The lattice size $N = 4 \times 4$. The thickness of links represents the magnitude of the doublon-holon correlation function. Color scale: Blue (Red) indicates the largest (smallest) link weights.

networks are substantially more robust to random attacks or failures than homogeneous networks [16, 17]. The robustness of the network from its heterogeneity seems to indicate a hidden symmetry in network space. More precisely, the symmetry describes a phenomenon that the network function and structure almost remain unchanged or invariant under random removal of its links. Thus it could allow us to define an order parameter in the Mott transition by using appropriate network measures.

The small-world network containing both strong clustering and shortest path length [18] has been considered to be a well-established fact in many real-world networks [19–22]. Recently the small-world idea has been successfully implemented in several statistical models as well [23]. In order to quantify the small-world network, we further propose the small-worldness as the order parameter indicating the hidden symmetry broken in network space, which would suddenly change while the Hubbard system undergoes a Mott transition. We find that it is able to extract the same Mott transition point obtained by conventional quantities, such as charge stiffness. Based on this result as well as previous evidences obtained from different models [23], we demonstrate that the hidden symmetry related to resilience of networks can provide us another route to identify various many-body phase transitions with/without Landau symmetry breakings.

II. MODEL AND METHOD

We begin with the 2D half-filled Hubbard model on a square lattice, which is relevant to correlated electron materials [24]:

$$\begin{aligned}\hat{H} &= \hat{H}_t + \hat{H}_U \\ &= -\sum_{\langle i,j\rangle,\sigma}\left(c^\dagger_{i\sigma}c_{j\sigma} + H.c.\right) + U\sum_i \hat{d}_i,\end{aligned} \tag{1}$$

where $\hat{d}_i = n_{i\uparrow}n_{i\downarrow}$ and $n_{i\sigma} = c^\dagger_{i\sigma}c_{i\sigma}$ with spin σ. In the following, we express the trial ground state in the finite system of N sites with (anti-)periodic boundary condition along the $\hat{x}$ ($\hat{y}$) direction to satisfy the closed-shell condition.

For our numerical simulations, we use the variational Monte Carlo (VMC) approach, which is based on the well-known Gutzwiller wave function [25]

$$|\Psi_G\rangle = \hat{P}_G|\Psi_{FL}\rangle, \tag{2}$$

where $|\Psi_{FL}\rangle$ is the Fermi liquid and $\hat{P}_G = g^{\sum_i \hat{d}_i}$. Here g is the parameter to adjust the number of doublons in the system. However, it is impossible to realize the nonmagnetic Mott transition by just using $|\Psi_G\rangle$. To demonstrate this transition, the simplest doublon-holon binding correlation must be considered [8, 26],

$$|\Psi_{DH}\rangle = \prod_i \left(1 - \mu\hat{Q}_i\right)|\Psi_G\rangle, \tag{3}$$

$$\hat{Q}_i = \hat{d}_i\prod_\delta\left(1-\hat{h}_{i+\delta}\right) + \hat{h}_i\prod_\delta\left(1-\hat{d}_{i+\delta}\right). \tag{4}$$

Here $\hat{h}_i = (1-n_{i\uparrow})(1-n_{i\downarrow})$ and δ runs over the nearest neighbors of the site i. μ is the parameter to control the number of isolated doublons and holons. If $\mu = 0$, $|\Psi_{DH}\rangle$ will be back to $|\Psi_G\rangle$ which is metallic for $U < \infty$. On the other side, $\mu = 1$, $|\Psi_{DH}\rangle$ becomes a completely doublon-holon bound state that seems to be insulating. It has been numerically proven that $|\Psi_{DH}\rangle$ is enough to describe the behavior of the Mott transition in VMC studies, even though we still need to consider more correlation factors to correctly capture the transition point [27]. Thus, it is reasonable to illustrate the occurrence of a nonmagnetic Mott transition by using the nearest-neighbor doublon-holon bound state.

In the language of network analysis, each network of N nodes can be described by the $N \times N$ adjacency matrix $\hat{A}$ [9]. A number of real systems, *e.g.* social networks, transportation networks, biological networks and so on, are better captured by weighted networks in which links use the weight to quantify their strengths. Here we consider lattice sites as nodes of the weighted network of which each weighted link between nodes i and j is expressed by the element of the adjacency matrix $\hat{A}_{ij}$. The link should carry the weight containing information about the relationship between electrons in the Hubbard system. According to the trial wave function we use, the weight of the network link is naively defined by the doublon-holon binding correlation. Namely, the lattice now breaks into N single sites called nodes, and then each pair of nodes i and j is reconnected by using the doublon-holon correlation function given by $\langle\hat{d}_i\hat{h}_j\rangle$.

Many of real networks have the property of relatively short average path length that is a shortest route running along the links of a network. In other words, most nodes may not be neighbors but can reach each other by a small number of steps. This is called small-world phenomena [28]. However, the Erdös-Rényi random networks showing the small-world effect fail to reproduce some important features of real networks, e.g. clustering. A small-world network including not only strong clustering but also short path length has thus been introduced to describe real networks by Watts and Strogatz [18]. Instead of the weighted clustering coefficient $\langle c\rangle$ [29] and the average path length $\langle d\rangle$ [30] commonly used in network analysis, an alternative measurement of the small-world property called "small-worldness" has been recently proposed [31]. The definition is based on the maximal tradeoff between high clustering and short path length. We can further define small-worldness as

$$\langle s\rangle \equiv \frac{\langle c\rangle}{\langle d\rangle}. \tag{5}$$

One can see more details in Ref. [23]. A network with larger $\langle s\rangle$ has a higher small-world level. Since the small-worldness simultaneously contains the information about both locality (weighted clustering coefficient $\langle c\rangle$) and non-locality (average path length $\langle d\rangle$) in network space, we can expect that the small-worldness is very suitable for describing the universal critical properties, especially the phase transitions without local order parameters. In fact, we have shown that the small-worldness indeed can behave like an order parameter in either continuous or topological phase transitions in our previ-



FIG. 2: (a) The kinetic energy E_t and (b) the interaction energy E_U/U of the 2D Hubbard Hamiltonian as a function of U for the lattice size $N = 16 \times 16$. (c) The charge structure factor N_k and (d) the momentum distribution function n_k along the nodal direction from M (π, π) to Γ $(0, 0)$ for different U.



FIG. 3: The doublon-holon correlation function $C_{dh}(R)$ of the 2D Hubbard model of lattice size $N = 16\times16$ vs distances R for various U.

ous works [23].

III. MOTT TRANSITION OF PHYSICAL OBSERVABLES

We now recall some facts about the nonmagnetic Mott transition arising from the doublon-holon binding effect [27]. The short-range doublon-holon binding correlation to describe the essence of the first-order Mott transition has been confirmed by several VMC calculations [32, 33]. A recent VMC study also concludes that the size of the doublon-holon bound state in the insulating phase is not beyond the next nearest neighbors [27]. This result thus motivates us to simply consider the nearest-neighbor doublon-holon binding correlation in the trial wave function $|\Psi_{DH}\rangle$. However, the critical issue is that the proposals on the order parameter are still controversial even if the metal-insulator transition is well defined. In the following, we therefore examine several conventional physical observables to detect the critical point.

Figure 2(a) and (b) show the kinetic energy $E_t(\equiv \langle \hat{H}_t \rangle)$ and the interaction energy $E_U/U(\equiv \langle \hat{H}_U \rangle/U)$ which is substantially the doublon density. It is obvious that both the kinetic energy and the doublon density have a discontinuity at the critical point U_c that has been indicated by the other VMC approach ($U_c = 8.575$) [27]. This is a typical sign of a first-order transition. Nevertheless, we cannot do anything but further compute the maximum decreasing (minimum increasing) rate of the interaction energy $-\frac{\partial E_U}{\partial U}$ (the kinetic energy $\frac{\partial E_t}{\partial U}$) if we want to precisely know the value of U_c.

To estimate the critical point more accurately, we may need to consider other physical quantities. First, we start the charge density correlation function in momentum space called charge structure factor,

$$N_k = \frac{1}{N} \sum_{i,j} \langle n_i n_j \rangle e^{ik\cdot(R_i - R_j)} - n^2, \tag{6}$$

where $n_i = \sum_\sigma n_{i\sigma}$ and n electron density. One can see that the important information about the metal-insulator transition is concealed in the charge structure factor N_k. Within the variational theory, $N_k \propto |k|$ for $|k| \to 0$ if there is no charge gap, whereas $N_k \propto |k|^2$ if a charge gap opens. As illustrated in Fig.2(c), the momentum dependence of N_k near the Γ point abruptly changes between $U = 8$ and 9, which coincides with the critical region determined above. Even so, the way to distinguish the insulating phase from the metallic phase will need much more efforts on the interpolation between lattice points.

Then, we consider the momentum distribution function, $n_k = \sum_\sigma \langle c^\dagger_{k\sigma} c_{k\sigma} \rangle$. According to Fermi liquid theory, the jump of n_k near the Fermi surface, namely, the quasiparticle renormalization factor, can be used to identify the metal-insulator transition. It is also in connection with the charge stiffness. We show a clear discontinuity of n_k around the Fermi surface for small U in Fig.2(d), thus suggesting that it is the metallic phase. As further increasing U, the jump is evidently reduced and remains very small for large $U(> 9)$. The small residual value of the jump is basically due to the finite size effect. That means we need to make more endeavors to indicate the critical point U_c estimated from the extrapolation to where the jump vanishes. So far, these observables we discuss above still cannot be thought of as the order parameter in the Mott transition even though they are able to distinguish the metallic and insulating phases.

IV. NETWORK ANALYSIS

In general, if a system shows the conventional long-range order, like ferromagnetism or superconductivity, the corresponding correlation function will be nonzero at its tail that can be used to define the order parameter. Otherwise, the correlation function would decay to zero at long distance for a

system does not break any symmetry. Here we would conjecture that it is impossible to find the order parameter derived from the doublon-holon correlation function because of unbroken symmetry in the nonmagnetic Mott transition. In what follows we shall illustrate a symmetry breaking unseen in real space.

From the doublon-holon binding effect leading to the nonmagnetic Mott transition we can reasonably infer that the correlation between a doublon and a holon provides a sufficient amount of data about the critical point U_c. We define the doublon-holon correlation function as follows,

$$C_{dh}(R) = \frac{1}{N}\sum_i \langle \hat{d}_i \hat{h}_{i+R} \rangle. \quad (7)$$

In Fig.3, we show how the doublon-holon correlation function changes as the Mott transition occurs. For $U = 0$, the Fermi-liquid ground state would not present unusual relationship between a doublon and a holon. As increasing U, a doublon starts to attract one holon at the nearest-neighbor site so that the doublon-holon correlation function slowly enhances only at $R = 1$. When $U > U_c$, $C_{dh}(R)$ rapidly increases at $R = 1$ as well as $C_{dh}(R = 1) \gg C_{dh}(R \neq 1)$ since the doublon-holon bound state begins to develop in the insulating phase. A key observation from Fig.3 is that while the system enters the insulating regime, the short-range parts ($R < 2$) of the doublon-holon correlation function immediately decay with distances but the long-range parts still remain similar to the metallic phase. In order to search a possible order parameter in network space, we will take advantage of this property of which the correlation function behaves differently for small and large U.

We now investigate how the network techniques perform in the face of the metal-insulator transition. Based on the observation on the doublon-holon correlation, a straightforward definition for the elements of the adjacency matrix $\hat{A}_{ij}$ is the magnitude of the normalized doublon-holon correlation function between the lattice sites i and j,

$$\hat{A}_{ij} = \frac{|\langle \hat{d}_i \hat{h}_j \rangle|}{max(|\langle \hat{d}_i \hat{h}_j \rangle|)}. \quad (8)$$

The complex topology of the weighted network extracted from $\hat{A}_{ij}$ has been shown in Fig.1 for the small lattice. Obviously, the weight of network links exhibits different distributions for the metallic and insulating phases even though all nodes in the network are completely connected. It would be just a trivial complete network if the network were binary. Next, we need to collect useful information from a large volume of network data.

In our previous studies [23], we have confirmed that the small-worldness $\langle s \rangle$ defined in Eq.(5) has the ability to detect the second-order and topological phase transitions. Here we further examine the feasibility of the small-worldness in the first-order phase transition without any symmetry breaking. In Fig.4(a), we illustrate the critical behavior of the small-worldness in the 2D Hubbard model. One can see that the $U-$dependence of the small-worldness behaves like an order parameter and drops down to almost zero near the critical



FIG. 4: (a) Small-worldness $\langle s \rangle$ of the 2D Hubbard model as a function of U for $N = 16 \times 16$. The inset shows the magnification near the critical point and includes the data points in a larger lattice system ($N = 20 \times 20$). (b) The probability distribution of the link weight W for different U. The bin size is chosen as 0.01 for clear demonstrations. The lattice size $N = 16 \times 16$.

point U_c in the finite system of size $N = 16 \times 16$. In the inset of Fig.4(a), we find that the critical point is indicated by the sudden reduction of $\langle s \rangle$ at U_c, in agreement with those results obtained above and in Ref. [27]. Notice that in a larger lattice of size $N = 20 \times 20$ the small-worldness for $U > U_c(\sim 8.9)$ is further reduced to zero. However, it is rather difficult to estimate U_c by using finite-size calculations since the Mott critical value of the doublon-holon binding state would become larger as increasing lattice size. In any case, this agreement convinces us that the small-worldness can be used as another quantity to characterize the nonmagnetic Mott transition instead of the charge stiffness.

To understand this result, we need to show how the weight distribution of network links enables the complex weighted network to illustrate the Mott transition. Fig.4(b) shows that the weights of network links display two bounded distributions with the asymmetric bimodal shape. The bimodal shape is an artifact, directly associated with the broken C_4 rotational symmetry by anti-periodic boundary condition along the $\hat{y}$ direction. At $U = 0$, few links have the strongest weight ($W \sim 1$) from which the weights for most links are not far away ($W \sim 0.75$). Thus, if we were to ignore the bimodal shape for the time being, the weighted network for $U = 0$ would exhibit the homogeneous distribution around $W = 1$, corresponding to the strongest small-world network whose most links resemble the highway in road networks.

As further increasing U, the weights for most links begin to move away from the largest weight ($W = 1$) and hence the distributions become more heterogeneous, leading to the reduction of the small-worldness as shown in Fig.4(a). When $U > U_c$, most links have very small weights ($W \sim 0$) and few links with the stronger weights ($W \sim 0.9$) show the broad distribution. The phenomena that most links belong to slow traffic lanes would result in almost zero small-worldness (weak clustering and long path). The same reasoning from the weight distribution of network links could be also applied to other many-body systems without local order parameters in real space. Hence this result strongly suggests that the small-worldness can be considered as a new order parameter in the network representation to capture the nonmagnetic Mott transition.

We have to mention a point now in passing. In the network

TABLE I: Examples of symmetry breakings and their corresponding order parameters.

Normal phase	Condensed phase	Broken symmetry	Order parameter
Paramagnet	Ferromagnet	Spin rotation	Magnetization
Metal	Superconductor	Global gauge	Pair condensate
Heterogeneous network	Homogeneous network	Network robustness	Small-worldness

representation, there exists a kind of hidden symmetry corresponding to the heterogeneous network with a broad weight distribution. This symmetry is related to the robustness of the network that is resilient to random attacks or failures [17]. The heterogeneity of network links implies that the weighted network is much more robust against attacks than the homogeneous networks, giving rise to the higher symmetry in network space. Conversely, the homogeneity of network links means that the weighted network becomes fragile to random attacks, and thus breaks the hidden symmetry. The hidden symmetry breaking is in analogy with typical examples of spontaneous symmetry breakings in condensed matter physics, such as the ferromagnet or superconductor (see the comparison in Table I). Therefore, we demonstrate that the small-worldness carrying the messages about both locality (clustering) and nonlocality (path length) is able to represent the order parameter describing the hidden symmetry broken in network space.

V. CONCLUSION

In this work we have addressed how to read useful information from the doublon-holon correlation function to identify the nonmagnetic Mott critical point by using complex network analysis. We have compared several conventional observables describing the Mott transition, and yet none of them can be formally treated as the order parameter. In the 2D Hubbard model, we have illustrated that one of many network measures, small-worldness, plays a significant role as a general order parameter concealed in network space. The critical point U_c extracted from the small-worldness is very close to the one obtained from other physical observables. Thus, it is anticipated that the small-worldness can be applied to the many-body systems beyond Landau symmetry-breaking theory.

The broadening of the weight distribution of network links across the critical point is responsible for the change of the small-worldness, which is analogous to the change of the speed limit on a road network from the highway to the slow lane. The phenomenon that the structure of the weighted network varies from heterogeneity to homogeneity implies a hidden symmetry broken−or, to put it another way, the reduction of the robustness to random attacks in the network space. We have confirmed that the broken symmetry can be successfully described by the small-worldness. The most important finding in this work is the hidden symmetry breaking in network space, namely, the reduction of network resilience, which has been listed in Table I.

The weight distribution of network links is able to uncover a wealth of complex topological information underneath correlation functions, and further comprehends the mechanism of the phase transition with/without any order parameter in real space. We propose that the small-worldness determined from various weighted networks could be a valuable tool to investigate quantum phase transitions. The finding has been corroborated by our previous results in the 1D quantum Ising and 2D classical XY models [23]. As a result, plus the evidence given in the 2D Hubbard model, one may conjecture that generally the point of view coming from complex network topology can be a vital component in studying the phase transitions in condensed matter physics.

Acknowledgments

We thank T.-K. Lee, F. Yang, and M.-C. Chang for helpful discussions. This work is supported by Chinese Academy of Engineering Physics and Ministry of Science and Technology. Most of calculations are performed in the National Center for High-performance Computing in Taiwan.

[1] N. F. Mott, Metal-Insulator Transitions Taylor & Francis, London (1974).
[2] M. Greiner *et al.*, Nature **415**, 39 (2002).
[3] T. Stöferle, H. Moritz, C. Schori, M. Kohl, and T. Esslinger, Phys. Rev. Lett. **92**, 130403 (2004).
[4] I. B. Spielman, W. D. Phillips, and J. V. Porto, Phys. Rev. Lett. **98**, 080404 (2007).
[5] M. Imada, A. Fujimori, and Y. Tokura, Rev. Mod. Phys. **70**, 1039 (1998).
[6] W. F. Brinkman and T. M. Rice, Phys. Rev. B **2**, 4302 (1970).
[7] M. A. Continentino, *Quantum Scaling in Many-Body Systems* (World Scientific) (2001).
[8] C. Castellani, C. Di Castro, D. Feinberg, and J. Ranninger, Phys. Rev. Lett. **43**, 1957 (1979).
[9] R. Albert and A.-L. Barabasi, Rev. Mod. Phys. **74**, 47 (2002).
[10] S. N. Dorogovtsev and J. F. F. Mendes, Adv. Phys. **51**, 1079 (2002).
[11] M. E. J. Newman, SIAM Rev. **45**, 167 (2003).
[12] S. Boccaletti *et al.*, Phys. Rep. **424**, 175 (2006).
[13] S. N. Dorogovtsev, A. V. Goltsev and J. F. F. Mendes, Rev. Mod. Phys. **80**, 1275 (2008).
[14] D. Grady, C. Thiemann, and D. Brockmann, Nature Communi-

cations **3** 864 (2012).
[15] T. Senthil, J. B. Marston, and M. P. A. Fisher, Phys. Rev. B **60**, 4245 (1999).
[16] R. Albert, H. Jeong, and A.-L. Barabasi, Nature **406**, 378 (2000).
[17] B. Wang, H. Tang, C. Guo, and Z. Xiu, Physica A, **363**, 591 (2006).
[18] D. J. Watts and S. H. Strogatz, Nature (London) **393**, 440 (1998).
[19] R. Albert, H. Jeong and A.-L. Barabasi, Nature **401**, 130 (1999).
[20] M. E. J. Newman, Proc. Natl. Acad. Sci. USA **98**, 404 (2001).
[21] P. S. Dodds, R. Muhamad and D. J. Watts, Science **301**, 827 (2003).
[22] L. Backstrom *et al.*, *Proceedings of the Third Annual ACM Web Science Conference* (WebSci 2012) pp. 33 (2012).
[23] C.-P. Chou and M.-C. Chang, arXiv:1308.0255 (2013).
[24] F. Gebhard, *The Mott Metal-Insulator Transition Models and Methods* (Springer) (1997).
[25] M. Gutzwiller, Phys. Rev. Lett. **10**, 159 (1963).
[26] H. Yokoyama, Prog. Theor. Phys. **108**, 59 (2002).
[27] T. Miyagawa and H. Yokoyama, J. Phys. Soc. Jpn. **80**, 084705 (2011).
[28] D. J. Watts, *Small worlds: the dynamics of networks between order and randomness*, Princeton University Press (1999).
[29] J. P. Onnela, J. Saramaki, J. Kertesz and K. Kaski, Phys. Rev. E **71**, 065103 (2005).
[30] M. E. J. Newman, *Networks: an introduction*, Oxford University Press (2010).
[31] M. D. Humphries and K. Gurney, PLoS ONE **3**, e0002051 (2008).
[32] H. Yokoyama and H. Shiba, J. Phys. Soc. Jpn. **59**, 3669 (1990).
[33] H. Yokoyama, Y. Tanaka, M. Ogata, and H. Tsuchiura, J. Phys. Soc. Jpn. **73**, 1119 (2004).